

**ISSN 0393-5965, Volume 26, Number 3**

Volume 26 · Number 3 · September 2010

ISSN 0393-5965

# Aerobiologia

International Journal of Aerobiology

 Springer

**This article was published in the above mentioned Springer issue.  
The material, including all portions thereof, is protected by copyright;  
all rights are held exclusively by Springer Science + Business Media.  
The material is for personal use only;  
commercial use is not permitted.  
Unauthorized reproduction, transfer and/or use  
may be a violation of criminal as well as civil law.**

# Assessment of bioaerosols at a concentrated dairy operation

Robert S. Dungan · April B. Leytem ·  
Sheryl A. Verwey · David L. Bjorneberg

Received: 31 July 2009 / Accepted: 12 January 2010 / Published online: 26 May 2010  
© US Government 2010

**Abstract** Increased bioaerosol loadings in downwind plumes from concentrated animal feeding operations (CAFOs) may increase the risk for allergy and infection in humans. In this study, we monitored airborne concentrations of culturable bacteria and fungi at upwind (background) and downwind sites at a 10,000 milking cow dairy over the course of a year. The average bacterial concentrations at the upwind site were  $8.4 \times 10^3$  colony forming units (CFU)  $\text{m}^{-3}$  and increased to  $9.9 \times 10^5$  CFU  $\text{m}^{-3}$  at the downwind edge of the cattle lots, decreasing to  $6.3 \times 10^4$  CFU  $\text{m}^{-3}$  200 m farther downwind. At the same sites, the average fungal concentrations were 515, 945, and 1,010 CFU  $\text{m}^{-3}$ , respectively. Significant correlations between the ambient weather conditions and airborne fungal and bacterial concentrations were identified. Sequence analysis of PCR-amplified DNA from bacterial clones and fungal isolates revealed genus and species level differences between upwind and downwind sites. Although we could not cultivate gram-negative bacteria, bacterial clones at downwind sites identified as being gram-negative matched with the following genera: *Acinetobacter*, *Bradyrhizobium*, *Escherichia*, *Idiomarina*, *Methylobacterium*,

*Ralstonia*, and *Novosphingobium*. Fungal isolates from downwind matched with the following genera: *Acremonium*, *Alternaria*, *Ascomycte*, *Aspergillus*, *Basidiomycete*, *Cladosporium*, *Davidiella*, *Doratomyces*, *Emericella*, *Lewia*, *Onygenales*, *Penicillium*, *Rhizopus*, and *Ulocladium*. None of the bacterial and fungal sequence matches were affiliated with genera and species known to be pathogenic to humans. Overall, the data suggest that exposure to bioaerosols in the downwind environment decreases with increasing distance from the open-lot dairy.

**Keywords** Airborne bacteria and fungi · Bioaerosols · CAFOs · Dairy · Impaction · Impingement · Polymerase chain reaction

## 1 Introduction

As populations increase and land resources decline, the intensification and industrialization of animal production facilities is projected to increase globally. In the United States of America, concentrated animal feeding operations (CAFOs) have been growing at a rapid pace since the 1960s (Centner 2003). Along with this growth, there is increasing concern regarding the effects of CAFOs on the health of workers, residents in nearby communities, livestock, and the environment. High animal densities are associated with the buildup of the overall microbial load in the

R. S. Dungan (✉) · A. B. Leytem · S. A. Verwey ·  
D. L. Bjorneberg  
Northwest Irrigation and Soils Research Laboratory,  
Agricultural Research Service, US Department  
of Agriculture, 3793 North 3600 East, Kimberly,  
ID 83341, USA  
e-mail: robert.dungan@ars.usda.gov

production environment by virtue of increased volumes of animals, feed, manure, and wastewater (Millner 2009). As a result, there is a concurrent increase in the bioaerosol (e.g., bacteria, fungi, virus, endotoxin) and dust load released from CAFOs (Sweeten et al. 1988; Wilson et al. 2002; Rule et al. 2005; Dungan and Leytem 2009a).

In Idaho, there are approximately 750 dairies and as many as 549,000 milking cows (USDA, National Agricultural Statistics Service 2008). Over the last decade, the milking cow population has increased by 88% and the trend is increasing toward larger concentrated facilities. While 46% of the dairy operations in Idaho contain <200 cows, some of the largest operations have as many as 10,000 animals. Due to the increased bacterial loading at concentrated dairy and other animal operations, there is a potential for offsite transport of bioaerosols and subsequent adverse health effects in humans and livestock (Wilson et al. 2002; Green et al. 2006; Schulze et al. 2006; Heederik et al. 2007). The inhalation of airborne microorganisms and their fragments can be detrimental to health through infection, toxicosis, or allergy (Crook and Sherwood-Higham 1997).

A large open-lot dairy in southern Idaho was targeted for this study. The specific objectives were to (1) measure the concentration of airborne microorganisms


at upwind (background) and downwind sites using all-glass impingers and single-stage cascade impactors; (2) determine statistical differences between airborne microorganism concentrations at each sample site and identify correlations with ambient weather data; (3) use a high-volume cyclonic sampler to capture airborne bacteria, then prepare clones from isolated DNA using PCR-amplified regions of 16S ribosomal DNA (rDNA) for sequence analysis and identification; and (4) PCR amplify the internal transcribed spacer (ITS) region of fungal isolates for sequence analysis and identification.

## 2 Materials and methods

### 2.1 Open-lot dairy

The dairy used in this study was a privately owned commercial dairy in southern Idaho, in a rural location, with 10,000 milking cows and a stocking density of approximately  $55 \text{ m}^2 \text{ cow}^{-1}$  (Fig. 1). This dairy is similar in configuration to most open-lot production facilities in southern Idaho. The operation consists of 24 open-lot pens, two milking parlors, hospital barn, maternity barn, manure solid separator, lagoon (liquid storage pond), and compost yard. There are approximately 13,000 cows on the facility

**Fig. 1** Map of the open-lot dairy with sampling sites



including milking cows, dry cows, and replacement heifers. Manure is scraped or vacuumed from feed alleys daily and placed into cells near the solid separator. The open-lot pens are harrowed daily when dry. The facility is surrounded by irrigated crop land on three sides and open range to the north. Three sites were used for sampling: an upwind site (approximately 200 m upwind from the open lots), 5 m downwind from the edge of lots (DW1), and 200 m downwind from the edge of the lots (DW2).

## 2.2 All-glass impingers

All-glass liquid impingers (SKC Inc., Eighty-Four, PA, USA) were utilized to capture airborne bacteria (i.e., heterotrophs, gram-negatives, *Escherichia coli*, and enterococci) and coliphage. Prior to their use, each impinger was autoclaved (30 min, 121°C, 1.23 atm) and then filled with 30-mL of sterile impingement solution (0.1% peptone, 0.1% antifoam B emulsion, pH 6.8). The impingers were mounted on tripods at a height and spacing distance of 1.5 m. The flowrate through each of the impingers was set to 8.5 L min<sup>-1</sup> using a Vac-U-Go sampling pump (SKC Inc.). While the manufacturer recommends a flowrate of 12.5 L min<sup>-1</sup>, we conducted a field study to test the recovery of heterotrophic bacteria. Our results demonstrated that bacterial recoveries at 8.5 versus 12.5 L min<sup>-1</sup> were, on average, 36% greater (unpublished data).

Bioaerosol samples were collected twice each day between the hours of 0800–1200 and 1300–1700. Three impingers were utilized at each sampling site and all samples were collected simultaneously. Samples were collected for 1 h and flowrate through the impingers was checked before and during operation with a rotameter. To reduce light penetration into the impingement solution, a plastic sleeve coated with aluminum foil was placed on the bottom portion of each impinger. After sampling, the collection vessels from the bottom of the impingers were aseptically stored and transported in a cooler with ice packs. Upon receipt at the laboratory, the samples were stored at 5°C for no longer than 18 h before being processed. The impingers were disinfected between sampling events by rinsing with 70% ethanol.

To account for any evaporation, the volume in each collection vessel was brought back up to 30 mL through additions of sterile impingement solution. The impingement solution was then serially diluted in

phosphate-buffered saline and 0.1-mL aliquots of the dilutions were spread directly onto Petri plates containing tryptic soy agar (TSA) and MacConkey agar. Cyclohexamide at 50 mg L<sup>-1</sup> was added to the TSA plates to inhibit the growth of fungi. The TSA and MacConkey agar were used for the enumeration of total heterotrophic and gram-negative bacteria, respectively. Through 47-mm, 0.45- $\mu$ m pore-size membrane filters (Pall Corp., Ann Arbor, MI, USA), 100-mL aliquots of the dilutions were filtered, which were then placed onto MI agar and *m*Enterococcus plates. The MI agar (US Environmental Protection Agency 2002; method 1604) and *m*Enterococcus agar (Eaton et al. 2005, method 9230C) were used for the detection and enumeration of total coliforms and *E. coli* and fecal streptococci, respectively. The agar plates were incubated aerobically under the following conditions: TSA, 25°C for 5 days; MacConkey, 35°C for 5 days; MI, 35°C for 1 day; and *m*Enterococcus, 35°C for 2 days. Coliphage were enumerated using the pour plate method (9224B) as described by Eaton et al. (2005). To ensure quality control, blanks, trip blanks, and positive controls were utilized for all microbiological assays.

## 2.3 Single-stage cascade impactors

Airborne fungi were collected directly on potato dextrose agar (PDA) through the use of single-stage cascade impactors (BioStage 200, SKC Inc.). The cascade impactors are equipped with 200 holes (0.25 mm diameter), and a QuickTake 15 sampling pump (SKC Inc.) was used to pull vacuum at a flowrate of 14 L min<sup>-1</sup>. The impactors were mounted 1.5 m off the ground at each of the sampling sites. Since only one impactor was available for each site, triplicate samples were collected in succession for a total of 2 or 5 min for each PDA plate. Fungal samples were collected twice each day, which occurred during the same times as the impinger samples were collected. The PDA plates were treated with 200 mg streptomycin sulfate L<sup>-1</sup> to inhibit bacterial growth. After 5 days of growth at 25°C, fungal colonies were enumerated.

A microscopic examination of spores and hyphae and sequencing of the ITS region was used to identify fungal isolates. Fungal isolates from the PDA plates were cultured for up to 2 weeks in potato dextrose broth at room temperature with shaking (50 rev min<sup>-1</sup>). The

isolates were then subsequently freeze dried and the DNA was extracted using a DNeasy Plant Mini Kit (Qiagen Inc., Valencia, CA, USA) according to the manufacturer's recommendations. Polymerase chain reaction (PCR) of ITS region was performed using the ITS1, ITS4, and ITS5 primers (White et al. 1990; Gardes and Bruns 1993) at a final concentration of 0.3  $\mu\text{M}$ . The PCR thermocycler conditions were 95°C for 10 min, then 30 cycles of 94°C for 30 s, 55°C for 40 s, and 72°C for 50 s, and single final extension at 72°C for 5 min. Products were visualized on 2.0% agarose gels by UV illumination and ethidium bromide staining.

#### 2.4 High-volume sampler

Samples were collected at sites upwind and downwind of the lots, wastewater lagoon, and compost piles using a SASS 2300 (Smart Air Sampler System, Research International, Monroe, WA, USA). The SASS 2300 is a cyclonic system designed to collect particles with an aerodynamic diameter of 1–10  $\mu\text{m}$ . Since only one SASS unit was available, samples at each site were collected at least one day apart. The system was operated at a flowrate of 325 L  $\text{min}^{-1}$  and sterile deionized (DI) water was used as the collection liquid as recommended by the manufacturer. The unit was placed approximately 1 m off the ground and operated for 8 h. Afterward, the concentrated liquid sample was transferred to a sterile 50-mL conical tube; the cyclone was subsequently rinsed twice with sterile DI (by temporarily operating the system for 1 min) and the rinsate was also added to the sterile tube. The SASS sample was transferred to the laboratory in a cooler and stored at  $-20^\circ\text{C}$  until processed. To process the samples, they were initially thawed at room temperature and then passed through a 0.4- $\mu\text{m}$  polycarbonate track-etch membrane. The membrane was added to a bead beating tube from an UltraClean soil DNA isolation kit (MoBio Laboratories, Inc., Carlsbad, CA, USA), rapidly vortexed for 1 min, and then the membrane was aseptically removed using sterile forceps. The tubes were subjected to bead beating for 2 min and processed according to the manufacturer's protocol.

Nested PCR of the DNA was performed by first using the primer pair BA8F and UN1492R (Reysenback et al. 1994) with the following program: 95°C for 5 min, then 30 cycles of 94°C for 45 s, 46°C for 30 s, and 72°C for 1 min, and a single final extension

at 72°C for 5 min. Then, the V1–V3 region of the 16S rDNA was amplified with the primer pair 63F and BA518R (Muyzer et al. 1993; Marchesi et al. 1998) with the following program: 95°C for 5 min, then 30 cycles of 92°C for 1 min, 55°C for 30 s, and 72°C for 1 min, and a single final extension at 72°C for 5 min. The nested PCR product from primer pair 63F-BA518R was cloned into pGEM-T Easy Vector (Promega Corp., Madison, WI, USA). White colonies were randomly selected, grown overnight at 37°C in Luria–Bertani medium containing 0.1 g ampicillin  $\text{L}^{-1}$ . Plasmids from *E. coli* JM109 transformants were isolated using the UltraClean mini plasmid prep kit (MoBio Laboratories, Inc.).

#### 2.5 Sequencing and identification

The plasmid inserts were amplified with the 63F primer and PCR product from the fungal isolates were sequenced by TACGen (Richmond, CA, USA) using an ABI 3730xl DNA Analyzer (Applied Biosystems, Foster City, CA, USA). Sequence identification was performed using the BLAST database at the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov>). The 16S rDNA sequences and representative sequences from the BLAST database were aligned with the BioEdit Sequence Alignment Editor (Hall 1999), and the aligned sequences were classified using the Greengenes Web site (<http://greengenes.lbl.gov/cgi-bin/nph-index.cgi>).

#### 2.6 Statistical analysis

All data were tested for normality using the Shapiro–Wilk test with the PROC CAPABILITY procedure of SAS (SAS Institute 2004). Data that were not normally distributed was log transformed prior to analysis with untransformed numbers presented in the text. The data were analyzed using the Mixed Procedure of SAS with date as the repeated measure and site as the subject. Means separation was carried out using the difference of the least squares means with the Tukey–Kramer adjustment and  $\alpha$  of 0.05. To determine the relationship between ambient weather data and airborne microorganism concentrations, Pearson correlation coefficients ( $r$ ) were calculated. Statements of statistical significance were based upon  $P < 0.05$  unless otherwise stated.

### 3 Results

The ambient weather conditions at the dairy are presented in Table 1. The weather data are averages from the periods during which bioaerosols were collected. The ambient temperature ranged from  $-6.2$  to  $30.2^{\circ}\text{C}$  with an average of  $14^{\circ}\text{C}$  during the study. The relative humidity ranged from 18 to 90% with an average of 48%. Solar radiation was generally the highest in April–September and lowest in October–March with an average of  $436\text{ W m}^{-2}$ . The wind speed

ranged from  $1.0$  to  $7.4\text{ m s}^{-1}$ , and the prevailing wind was from the west. Bioaerosols were only collected when the wind was predominantly from the west (avg. =  $244^{\circ}$ , SEM =  $7.4^{\circ}$ ) or east (avg. =  $115^{\circ}$ , SEM =  $13.6^{\circ}$ ). Bioaerosol sampling was not initiated during precipitation events; however, in two instances, light rain started once sampling had commenced. The upwind and downwind sampling sites at the open-lot dairy are shown in Fig. 1.


All-glass impinger samples were collected during all months, except December, January, and February,

**Table 1** Average ambient weather data and lot conditions during sample collection

| Year        | Day-month    | Air temperature ( $^{\circ}\text{C}$ ) | RH (%) | Solar Radiation ( $\text{W m}^{-2}$ ) | WS ( $\text{m s}^{-1}$ )                    | Lot conditions   |
|-------------|--------------|--|--------|---------------------------------------|---|--|
| 2008        | 21-April     | 6.4                                    | 29     | 712                                   | 2.8   | Lots very wet with standing water in several lots along eastern edge, manure piles present |
|             | 28-April     | 17.1                                   | 25     | 616                                   | 3.5   |  |
|             | 19-May       | 24.2                                   | 34     | 702                                   | 4.2   | All lots were cleaned out and manure piles removed, lots were dry                          |
|             | 22-May       | 10.7                                   | 57     | 344                                   | 2.4   |  |
|             | 16-June      | 24.8                                   | 27     | 695                                   | 2.2   | Lots were dry, some buildup of manure piles  |
|             | 18-June      | 20.9                                   | 38     | 679                                   | 3.2   |  |
|             | 28-July      | 24.5                                   | 43     | 652                                   | 2.1   | Lots were dry, manure piles were present, new soil added to lots on eastern side           |
|             | 30-July      | 20.3                                   | 26     | na                                    | 3.7   |  |
|             | 18-August    | 30.2                                   | 18     | 611                                   | 3.5   | Lots were dry, some buildup of manure piles  |
|             | 20-August    | 25.0                                   | 27     | 587                                   | 1.0   |  |
|             | 22-September | 13.3                                   | 60     | 458                                   | 4.1   | Lots were dry, some buildup of manure piles  |
|             | 25-September | 19.6                                   | 28     | 288                                   | 1.0   |  |
|             | 27-October   | 16.0                                   | 19     | 302                                   | 2.1   | Lots were wet and some areas very muddy, manure piles present                              |
|             | 29-October   | 11.2                                   | 47     | 279                                   | 1.6   |  |
|             | 17-November  | 10.5                                   | 48     | 228                                   | 2.7   | Lots were wet and some areas very muddy, manure piles present                              |
| 19-November | 5.3          | 69                                     | 175    | 2.7                                   |   |  |
| 15-December | $-6.2$       | 86                                     | 118    | 7.4                                   | Lots were frozen over, manure piles present |  |
| 2009        | 27-January   | 12.3                                   | 82     | na                                    | 1.5   | Lots were frozen over, manure piles present  |
|             | 29-January   | 0.3                                    | 73     | 385                                   | 4.1   |  |
|             | 23-February  | 4.8                                    | 90     | 332                                   | 3.4   | Lots were mostly frozen, some thawing with standing water in places, manure piles present  |
|             | 25-February  | 5.6                                    | 71     | 250                                   | 2.4   |  |
|             | 16-March     | 12.0                                   | 55     | 136                                   | 3.8   | Lots were very wet, with standing water and mud in places, manure piles present            |
|             | 19-March     | 12.5                                   | 44     | 597                                   | 1.9   |  |

na not available,  
RH relative humidity, WS  
wind speed

as the impingement solution was prone to freezing. Therefore, impinger samples were only collected on 17 of 24 possible days. No culturable gram-negatives and coliphage were recovered during the course of the study. While the samples were occasionally positive for *E. coli* and enterococci at the downwind edge of the lot, the concentrations were relatively low. *Escherichia coli* was recovered on 4 separate events at concentrations ranging from 8 to 67 CFU m<sup>-3</sup>; enterococci was recovered on nine separate events at concentrations ranging from 2 to 21 CFU m<sup>-3</sup> (data not shown). Airborne heterotroph concentrations were highly variable during the study, resulting in high standard errors at each measurement time (Table 2). At the upwind site, the heterotroph concentrations ranged from  $9.1 \times 10^2$  to  $3.7 \times 10^4$  CFU m<sup>-3</sup>. At DW1, the heterotroph concentrations ranged from  $1.5 \times 10^4$  to  $1.4 \times 10^7$  CFU m<sup>-3</sup> and were always greater than upwind concentrations. At DW2, the concentrations ranged from  $4.4 \times 10^3$  to  $5.3 \times 10^5$  CFU m<sup>-3</sup> and were generally lower than DW1, but greater than upwind concentrations. Heterotroph concentrations tended to be the most consistent at the upwind site during July through October. An analysis of variance was performed on the data to determine the effect of sample site on the airborne heterotroph concentration. The average heterotroph concentrations were  $8.4 \times 10^3$ ,  $9.9 \times 10^5$ ,



**Fig. 2** Average airborne heterotroph concentrations measured at the upwind and downwind sites on the open-lot dairy. Letters within the columns indicate significant differences between the sample sites ( $P < 0.05$ ). Downwind 1, 5 m from the edge of the lot; Downwind 2, 200 m from edge of lot

and  $6.3 \times 10^4$  CFU m<sup>-3</sup> for the upwind, DW1, and DW2 sites, respectively (Fig. 2). The effect of site was only significant between the upwind and DW1 sites ( $P = 0.046$ ), but not between the two downwind sites or the upwind and DW2 site ( $P > 0.08$ ). There were no statistically significant correlations between air temperature, relative humidity, or solar radiation and airborne heterotroph concentration at each sample site

**Table 2** Average airborne concentrations of heterotrophs at the open-lot dairy

| Year | Day-month    | Upwind (CFU m <sup>-3</sup> ) | Downwind 1 (CFU m <sup>-3</sup> ) | Downwind 2 (CFU m <sup>-3</sup> ) |
|------|--------------|-------------------------------|-----------------------------------|-----------------------------------|
| 2008 | 21-April     | 915 ± 621 <sup>a</sup>        | 20232 ± 2350                      | 6164 ± 3507                       |
|      | 28-April     | 5238 ± 3660                   | 14593 ± 2824                      | 6916 ± 1877                       |
|      | 19-May       | 9314 ± 5663                   | 14349132 ± 14181077               | 124302 ± 93406                    |
|      | 16-June      | 9319 ± 2187                   | 50629 ± 23314                     | 13960 ± 3690                      |
|      | 18-June      | 37571 ± 19156                 | 55963 ± 6943                      | 19946 ± 4022                      |
|      | 28-July      | 1750 ± 700                    | 100434 ± 37223                    | 9115 ± 2029                       |
|      | 30-July      | 2184 ± 809                    | 168856 ± 47350                    | 54745 ± 9858                      |
|      | 18-August    | 16176 ± 8615                  | 122475 ± 35954                    | 29799 ± 7121                      |
|      | 20-August    | 3725 ± 661                    | 24602 ± 7676                      | 7534 ± 2336                       |
|      | 22-September | 2190 ± 714                    | 511118 ± 128192                   | 529687 ± 314082                   |
|      | 25-September | 5597 ± 1782                   | 297077 ± 96683                    | 118400 ± 17724                    |
|      | 27-October   | 8373 ± 3970                   | 535564 ± 295209                   | 44719 ± 11013                     |
|      | 29-October   | 9668 ± 3003                   | 126224 ± 33178                    | 25012 ± 6135                      |
| 2009 | 17-November  | 11188 ± 5713                  | 104021 ± 43227                    | 32271 ± 4618                      |
|      | 19-November  | 4615 ± 2053                   | 37573 ± 5599                      | 19216 ± 7352                      |
|      | 16-March     | 3713 ± 1825                   | 56111 ± 24479                     | 30065 ± 10499                     |
|      | 19-March     | 11713 ± 4784                  | 191806 ± 79870                    | 4455 ± 1727                       |

<sup>a</sup> Standard error of the mean ( $n = 6$ )

( $P > 0.13$ ). However, there was a positive correlation between wind speed and heterotroph concentration at DW1 ( $r = 0.23$ ,  $P = 0.02$ ) and DW2 ( $r = 0.30$ ,  $P = 0.002$ ).

A more detailed analysis of the airborne bacterial community was conducted by collecting aerosol samples using a high-volume sampler. DNA extracts from the samples were subjected to PCR using

**Table 3** Sequence matches of bacterial clones from upwind and downwind of various locations

| Clone | Closest GenBank match                             | Source                           | % Match | Accession no. | Phylum         |
|-------|---|----------------------------------|---------|---------------|----------------|
| U-1   | <i>Kocuria</i> sp. L5                             | Kartchner Caverns, Arizona       | 95      | DQ192212.1    | Actinobacteria |
| U-2   | <i>Actinobacteria</i> clone DOK_DONFYM_clone811   | Environmental sample             | 94      | DQ829010.1    | Actinobacteria |
| U-3   | <i>Ralstonia pickettii</i> strain 149             | Water Purifiers                  | 99      | EU730922.1    | Proteobacteria |
| U-4   | Uncultured <i>Microvirga</i> sp. clone AUVE_01B06 | Cropland                         | 99      | EF650876.1    | Proteobacteria |
| U-5   | <i>Delftia</i> sp. CAT3-4-2                       | Wastewater treatment pond        | 99      | FJ594443.1    | Proteobacteria |
| U-6   | <i>Nocardioideae bacterium</i> Ellin5426          | Soil                             | 98      | AY673153.1    | Actinobacteria |
| U-7   | <i>Bradyrhizobium japonicum</i>                   | Ice core sample                  | 96      | AY169423.1    | Proteobacteria |
| U-8   | <i>Pelomonas aquatica</i> strain CCUG52631        | Industrial water                 | 98      | AM501437.1    | Proteobacteria |
| U-9   | <i>Delftia</i> sp. CAT3-4-2                       | Wastewater treatment pond        | 99      | FJ594443.1    | Proteobacteria |
| U-10  | <i>Roseomonas aquatica</i> type strain TR53T      | Drinking water                   | 97      | AM231587.1    | Proteobacteria |
| CL-1  | <i>Ruminococcaceae bacterium</i> clone EMP_N10    | Fecal sample                     | 98      | EU794265.1    | Firmicutes     |
| CL-2  | <i>Bacteroidales bacterium</i> clone Cow183       | Cow feces                        | 99      | AY859656.1    | Bacteroidetes  |
| CL-3  | <i>Firmicutes bacterium</i> clone B11             | Raw cow milk                     | 99      | EU029233.1    | Firmicutes     |
| CL-4  | <i>Bacteroidetes bacterium</i> clone 912-C9       |                                  | 86      | EU315425.1    | Bacteroidetes  |
| CL-5  | <i>Firmicutes bacterium</i> clone B11             | Raw cow milk                     | 99      | EU029233.1    | Firmicutes     |
| CL-6  | <i>Acinetobacter</i> sp. TS11                     | Arsenic-contaminated environment | 97      | EU073077.1    | Proteobacteria |
| CL-7  | <i>Corynebacterium</i> sp. BBDP21                 | Swine effluent holding pit       | 94      | DQ337522.1    | Actinobacteria |
| CL-8  | <i>Escherichia</i> sp. 4094                       | Swine                            | 100     | FJ405352.1    | Proteobacteria |
| CL-9  | Clostridiales Family XIII bacterium clone EMP_F2  | Fecal sample                     | 94      | EU794167.1    | Firmicutes     |
| L-1   | <i>Atopostipes suicloacalis</i> strain PPC79      | Swine feces                      | 95      | AF445248.2    | Firmicutes     |
| L-2   | <i>Bradyrhizobium</i> sp. SEMIA 929               | <i>Ornithopus sativus</i>        | 100     | FJ390938.1    | Proteobacteria |
| L-3   | <i>Methylobacterium</i> sp. clone 1P-1-I17        | Environmental                    | 95      | EU704833.1    | Proteobacteria |
| L-4   | <i>Cyanobacterium</i> clone 3                     | Environmental                    | 97      | FJ024312.1    | Cyanobacteria  |
| L-5   | <i>Clostridium disporicum</i> strain NML 05A027   | Clinical isolate                 | 98      | DQ855943.1    | Firmicutes     |
| L-6   | <i>Idiomarina</i> sp. C4                          |                                  | 97      | EF554872.1    | Proteobacteria |
| L-7   | <i>Oscillatoria</i> sp. clone AV_4R-S-A21         | Commercial aircraft cabin air    | 96      | EU341185.1    | Cyanobacteria  |
| L-8   | <i>Corynebacterium efficiens</i> YS-314           |                                  | 95      | BA000035.2    | Actinobacteria |
| L-9   | <i>Clostridium</i> sp. clone MS159A1_E02          | Human gastrointestinal           | 99      | EF705259.1    | Firmicutes     |
| C-1   | <i>Planococcus</i> sp. YIM C832                   | Haloalkaline soil                | 98      | EU135681.1    | Firmicutes     |
| C-2   | <i>Firmicutes bacterium</i> clone QEDP3DD01       | Mesophilic anaerobic digester    | 99      | CU924228.1    | Firmicutes     |
| C-3   | <i>Ralstonia pickettii</i> strain 149             | Water Purifiers                  | 100     | EU730922.1    | Proteobacteria |
| C-4   | <i>Flavobacteriaceae bacterium</i> YIM C421       | Haloalkaline soil                | 93      | EU135615.1    | Bacteroidetes  |
| C-5   | <i>Clostridium lituseburense</i> strain H17       | Anaerobic digester               | 99      | EU887828.1    | Firmicutes     |
| C-6   | <i>Clostridium</i> sp. clone M139                 | Raw cow milk                     | 94      | EU029303.1    | Firmicutes     |
| C-7   | <i>Novosphingobium</i> sp. YC6720                 |                                  | 98      | EU707558.1    | Proteobacteria |
| C-8   | <i>Bacteroidales bacterium</i> clone CG2          | Cow feces                        | 93      | EU573809.1    | Bacteroidetes  |

U upwind, CL downwind of cattle lots, L downwind of lagoon, C downwind of compost piles



universal bacterial primers to amplify a section of 16S rDNA approximately 450 bp in length. The PCR products were used to produce representative clones that were subsequently used for sequence analysis and identification (Table 3). Data are presented for a total of 36 clones which were 86–100% homologous with known bacterial sequences. However, only 24 of 36 isolates could be identified to the genus level, with the remaining identified at the family level and higher. The clones from the upwind site and downwind of the cattle lots, lagoon, and compost piles were affiliated with the following phyla: *Actinobacteria*, *Bacteroidetes* (4 clones), *Firmicutes* (11 clones), and *Proteobacteria* (14 clones from the  $\alpha$ -,  $\beta$ -, and  $\gamma$ -subdivision). Two bacterial clones were also identified as belonging to *Cyanobacteria*, but only downwind of the lagoon. At the upwind site, bacterial clones were affiliated with bacteria mostly from environmental origins and belonging to the following genera: *Bradyrhizobium*, *Delftia*, *Kocuria*, *Microvirga*, *Pelomonas*, *Rolstonia*, and *Roseomonas*. At the downwind sites, bacterial

clones were identified with the following genera: *Acinetobacter*, *Clostridium*, *Corynebacterium*, *Escherichia*, *Methylobacterium*, *Oscillatoria*, *Ralstonia*, and *Novosphingobium*. Of the 26 downwind clones, 10 matched sequences from isolates whose source was feces (cow, swine) and raw cow milk.


Total airborne fungal concentrations are presented in Table 4. At the upwind site, the fungal concentrations ranged from 36 to 1,289 CFU m<sup>-3</sup>. At DW1 and DW2, the respective concentrations ranged from 181 to 2,314 CFU m<sup>-3</sup> and 192 to 5,274 CFU m<sup>-3</sup>. The average concentrations at the upwind, DW1, and DW2 sites were 515, 945, and 1,010 CFU m<sup>-3</sup>, respectively (Fig. 3). To determine the effect of site on the airborne fungal concentrations, an analysis of variance was performed. The effect of site was determined to be significant between the upwind and downwind sites ( $P < 0.0001$ ), but not between the two downwind sites ( $P = 0.73$ ). At the upwind site, a correlation analysis between the airborne fungal concentration and ambient weather data confirmed that there was a significant

**Table 4** Average airborne concentrations of fungi at the open-lot dairy

| Year        | Day-month                | Upwind<br>(CFU m <sup>-3</sup> ) | Downwind 1<br>(CFU m <sup>-3</sup> ) | Downwind 2<br>(CFU m <sup>-3</sup> ) |
|-------------|--------------------------|----------------------------------|--------------------------------------|--------------------------------------|
| 2008        | 21-April                 | 272 ± 78 <sup>a</sup>            | 1229 ± 258                           | 1148 ± 308                           |
|             | 28-April                 | 156 ± 47                         | 181 ± 66                             | 192 ± 50                             |
|             | 19-May                   | 240 ± 62                         | 379 ± 69                             | 629 ± 69                             |
|             | 22-May <sup>b</sup>      | 143 ± 48                         | 140 ± 26                             | 751 ± 327                            |
|             | 16-June                  | 584 ± 145                        | 408 ± 94                             | 560 ± 115                            |
|             | 18-June                  | 987 ± 303                        | 521 ± 80                             | 346 ± 52                             |
|             | 28-July                  | 1289 ± 161                       | 499 ± 95                             | 864 ± 155                            |
|             | 30-July                  | 927 ± 149                        | 490 ± 86                             | 573 ± 88                             |
|             | 18-August                | 949 ± 73                         | 447 ± 42                             | 501 ± 87                             |
|             | 20-August                | 517 ± 109                        | 505 ± 77                             | 312 ± 59                             |
|             | 22-September             | 836 ± 192                        | 885 ± 215                            | 511 ± 69                             |
|             | 25-September             | 871 ± 228                        | 1458 ± 352                           | 1710 ± 364                           |
|             | 27-October               | 436 ± 57                         | 649 ± 89                             | 1173 ± 74                            |
|             | 29-October               | 813 ± 59                         | 1182 ± 159                           | 854 ± 67                             |
|             | 17-November              | 352 ± 45                         | 886 ± 120                            | 1048 ± 303                           |
| 19-November | 369 ± 57                 | 1523 ± 190                       | 749 ± 110                            |                                      |
| 15-December | 679 ± 174                | 1576 ± 403                       | 943 ± 128                            |                                      |
| 2009        | 27-January               | 36 ± 0                           | 1524 ± 180                           | 5274 ± 1935                          |
|             | 29-January               | 37 ± 8                           | 432 ± 89                             | 310 ± 127                            |
|             | 23-February <sup>b</sup> | 206 ± 31                         | 2314 ± 649                           | 1193 ± 287                           |
|             | 25-February              | 206 ± 49                         | 2295 ± 250                           | 1858 ± 374                           |
|             | 16-March                 | 777 ± 95                         | 1703 ± 183                           | 904 ± 169                            |
|             | 19-March                 | 157 ± 24                         | 516 ± 115                            | 829 ± 145                            |

<sup>a</sup> Standard error of the mean ( $n = 12$ )

<sup>b</sup> Light rain event occurred during sampling



**Fig. 3** Average airborne fungal concentrations measured at the upwind and downwind sites on the open-lot dairy. Letters within the columns indicate significant differences between the sample sites ( $P < 0.05$ ). Downwind 1, 5 m from the edge of the lot; Downwind 2, 200 m from edge of lot

effect of air temperature ( $r = 0.23$ ,  $P = 0.008$ ). At the DW1 site, there was a significant effect of air temperature ( $r = -0.42$ ,  $P < 0.0001$ ), relative humidity ( $r = 0.47$ ,  $P < 0.0001$ ), and solar radiation ( $r = -0.45$ ,  $P < 0.0001$ ), while at DW2 there was a significant effect of air temperature ( $r = -0.20$ ,  $P = 0.03$ ), relative humidity ( $r = 0.31$ ,  $P = 0.0005$ ), and solar radiation ( $r = -0.32$ ,  $P = 0.004$ ). Interestingly, at DW1, there was a slight negative correlation between wind speed and fungal concentration ( $r = -0.23$ ,  $P = 0.009$ ), which is in contrast to the positive correlation seen between wind speed and airborne heterotroph concentration.

To identify the predominant filamentous fungi present at the dairy, unique isolates were obtained from the impaction plates after each sampling event. A total of 13 isolates from upwind and 18 isolates from the downwind edge of the lots were obtained. The ITS region of the fungal isolates was PCR-amplified and the products were subjected to sequence analysis. The isolates were 78–100% homologous with known fungal sequences; all but one isolate could not be identified to the genus level (i.e., clone CL-7). Upwind isolates were identified as being from the following genera: *Actinomucor*, *Aspergillus*, *Aureobasidium*, *Epicoccus*, *Fusarium*, *Neosartorya*, *Penicillium*, *Phoma*, *Radulidium*, *Talaromyces*, and *Ulocladium* (Table 5). Downwind from the lots, the following genera were identified: *Acremonium*, *Alternaria*, *Ascomycte*, *Aspergillus*, *Basidiomycete*, *Cladosporium*, *Davidiella*,

*Doratomyces*, *Emericella*, *Lewia*, *Onygenales*, *Penicillium*, *Rhizopus*, and *Ulocladium* (Table 5).

#### 4 Discussion

Bioaerosols are introduced into the atmosphere from various agricultural, municipal, industrial, and natural sources (Blanchard and Syzdek 1970; Brandi et al. 2000; Wilson et al. 2002; Brooks et al. 2005; Fracchia et al. 2006; Green et al. 2006; Spaan et al. 2006; Millner 2009). Airborne microorganisms or their components are typically associated with particulate matter or surrounded by a thin layer of water and have an aerodynamic diameter range of 0.5–100  $\mu\text{m}$  (Lighthart 1994; Cox 1995a). Aerosol particles 1–5  $\mu\text{m}$  are of most concern as they are readily inhaled or swallowed (Stetzenbach 2007). When bioaerosols are released from a source, they are transported short and long distances in air currents and are eventually deposited on surfaces (Brown and Hovmoller 2002; Jones and Harrison 2004). The transport and deposition of bioaerosols is affected by their physical properties and environmental conditions they encounter while airborne. Meteorological conditions that significantly affect the transport of bioaerosols are wind velocity, thermal gradients, and magnitude of air currents (Cox 1995a; Mohr 2007; Stetzenbach 2007). Important physical properties are size, density, and shape of the particles, while deposition of bioaerosols occurs through gravitational settling, wind impaction onto surfaces, adhesion, and other mechanisms (Wickman 1994). Factors that affect the viability of airborne microorganisms have been extensively studied; these include temperature, relative humidity, and solar radiation (Poon 1966; Marthi et al. 1990; Cox 1995b; Mohr 2007). In general, studies have shown decreased viability with decreases in relative humidity and increases in temperature and solar radiation. Oxygen concentration is also known to affect microbial survival, as it is involved in the inactivation of bioaerosols through the production of oxygen free radicals. While the above-mentioned atmospheric conditions influence viability, microbial factors such as the type, species, and strain of an organism also affect its airborne survival (Songer 1967; Ehrlich et al. 1970).

Compared to the upwind site, the airborne concentration of heterotrophic bacteria were statistically

**Table 5** Sequence matches of fungal isolates from upwind and downwind of the cattle lots

| Clone | Closest GenBank match                            | Source                                     | % Match | Accession no. |
|-------|--|--|---------|---------------|
| U-1   | <i>Epicoccum</i> sp. G7A                         | <i>Alliaria petiolata</i> (garlic mustard) | 91      | EF432273.1    |
| U-2   | <i>Phoma medicaginis</i> var. <i>medicaginis</i> | <i>Medicago sativa</i> (alfalfa)           | 99      | DQ109960.1    |
| U-3   | <i>Fusarium tricinctum</i>                       | <i>Picea mariana</i> (black spruce)        | 99      | DQ132835.1    |
| U-4   | <i>Fusarium equiseti</i>                         | Sorghum                                    | 98      | EU595566.1    |
| U-5   | <i>Penicillium vinaceum</i> strain P11.5         | Vampire bat guano                          | 98      | EU833227.1    |
| U-6   | <i>Penicillium commune</i> strain P4.2           | Cave soil                                  | 75      | EU833216.1    |
| U-7   | <i>Ulocladium</i> sp.                            | <i>Centaurea stoebe</i> (endophyte)        | 99      | EF589899.1    |
| U-8   | <i>Neosartorya fischeri</i> isolate wb162        | Nasal mucus                                | 88      | AF455541.1    |
| U-9   | <i>Aspergillus niger</i> strain MPVCT 344        | Vineyard                                   | 99      | EU440778.1    |
| U-10  | <i>Actinomyces elegans</i>                       | Potato pulp                                | 99      | AB113008.1    |
| U-11  | <i>Talaromyces flavus</i> isolate hn-50-1        | Insect-associated isolate                  | 99      | EU287814.1    |
| U-12  | <i>Radulidium subulatum</i> strain CBS 287.84    | Incubator for cell cultures                | 98      | EU041789.1    |
| U-13  | <i>Aureobasidium pullulans</i> strain SUB 04-313 |  | 97      | FJ744598.1    |
| CL-1  | <i>Rhizopus stolonifer</i> strain ATCC 6227b     | ATCC isolate                               | 78      | AF543526.1    |
| CL-2  | <i>Aspergillus fumigatus</i> strain F3           | Rhizospheric soil                          | 99      | FJ214371.1    |
| CL-3  | <i>Aspergillus niger</i> strain CH-A2010         | Tequila industry waste                     | 99      | FJ668837.1    |
| CL-4  | <i>Lewia infectoria</i> isolate T3549            | <i>Medicago lupulina</i> (black medick)    | 99      | EF104194.1    |
| CL-5  | <i>Basidiomycete</i> sp. nasa47                  | Atacama desert, Chile                      | 97      | DQ683975.1    |
| CL-6  | <i>Arthopyreniaceae</i> IBL 03137                | <i>Coffea arabica</i>                      | 99      | DQ682563.1    |
| CL-7  | <i>Cladosporium cladosporioides</i>              | <i>Phragmites australis</i> (common reed)  | 98      | AJ300334.1    |
| CL-8  | <i>Doratomyces</i> sp. HZ-10                     | Forest soil                                | 99      | EU301641.1    |
| CL-9  | <i>Alternaria</i> sp. FTJZZJ02                   | <i>Fritillaria thunbergii</i> Miq.         | 99      | FJ196610.1    |
| CL-10 | <i>Penicillium soppii</i> strain IBT 19343       |  | 100     | AM901674.1    |
| CL-11 | <i>Emericella rugulosa</i> strain SRRC 1173      |  | 99      | EU289916.1    |
| CL-12 | <i>Onygenales</i> sp. BC8                        | Ross Sea region of Antarctica              | 99      | DQ317338.1    |
| CL-13 | <i>Davidiella macrospore</i> strain CBS 107.20   | Iris species                               | 99      | EF679369.1    |
| CL-14 | <i>Acremonium exuviarum</i> strain BMB4          | Indoor isolate                             | 100     | AY882946.1    |
| CL-15 | <i>Penicillium janthinellum</i> strain P11.16    | Vampire bat guano                          | 98      | EU833221.1    |
| CL-16 | <i>Ascomycete</i> sp. BC20                       | Ross Sea region of Antarctica              | 100     | DQ317353.1    |
| CL-17 | <i>Ulocladium</i> sp.                            | Ross Sea region of Antarctica              | 99      | DQ317352.1    |
| CL-18 | <i>Penicillium roqueforti</i> strain FSU 503     |  | 99      | EU484267.1    |

U upwind, CL downwind of cattle lots

higher at the downwind edge of the lot (Fig. 2). The fact that the concentrations were higher at the downwind site supports the premise that animal confinement tends to increase the overall microbial load. As wind moves across the open lots, it aerosolizes particulate matter and fungal spores, leading to increases in the bioaerosol concentration in the downwind plume. While wind is a dominant factor, cow activity and lot harrowing are other mechanisms by which particulate matter are aerosolized in the lots. As the downwind distance from the source increases, one can expect bioaerosol

concentrations to decrease as gravitational settling of particles occurs and other depositional mechanisms take place. Gravitational settling and impaction is an important transport mechanism by which suspended particles  $>5 \mu\text{m}$  in diameter are removed from the air (Nicholson 1995). For small particles ( $<0.1 \mu\text{m}$  in diameter), Brownian diffusion is the dominant transport mechanism, which increases with increasing temperature and decreasing particle size.

At 200 m from the edge of the lot (i.e., DW2), the heterotroph concentration was 16-fold lower than at 5 m from the lot edge (i.e., DW1), although not a

significant difference (Fig. 2). The lower concentration at DW2 could be a result of particles settling out, or a combination of other factors, such as loss of viability and/or dilution. In a study conducted by Green et al. (2006), total culturable bacteria were monitored up to 150 m downwind from a swine confinement operation. Inside the swine facility, the average airborne bacterial concentration was  $1.8 \times 10^3$  CFU m<sup>-3</sup>. The airborne bacterial concentration was shown to steadily decrease with distance from the swine operation, with concentrations at 150 m downwind being only 2.5-fold higher than at an upwind site at 82 CFU m<sup>-3</sup>. In our study, the airborne heterotroph concentration was 8-fold higher at DW2 than the upwind site, but it was determined that these were statistically similar concentrations. Research on the land application of biosolids or at composting facilities has shown that bioaerosol concentrations decrease with increasing distance from the source (Brooks et al. 2004; Taha et al. 2005). Similarly, decreases in bioaerosol concentrations with increasing distance from wastewater treatment plants and land application of wastewater have been demonstrated (Brooks et al. 2004; Millner 2009). However, in the latter case, the microorganisms are associated with fine water droplets, but in our study they are likely bound to solid particles which certainly influence their viability. Desiccation of droplets is the main factor responsible for inactivation of bioaerosols in liquids (Mohr 2007).

A variety of airborne gram-negative bacteria, belonging to the families *Enterobacteriaceae*, *Pseudomonadaceae*, and *Neisseriaceae*, have been recovered in calf and other animal houses (Zucker et al. 2000; Millner 2009). In cow and calf stable,  $\leq 6.5\%$  of the viable airborne aerobic bacteria were identified as being gram-negative (Zucker and Müller 1998). A similar percentage of the airborne bacterial population was also determined to be gram-negative in swine houses (Chang et al. 2001; Kiekhaefer et al. 1995). While we did recover *E. coli* at relatively low concentrations at the downwind edge of the lot, we were unsuccessful in cultivating gram-negatives on selective agar. It is possible that the airborne gram-negative populations were not highly concentrated and, in combination with a 1 h sampling time with the all-glass impinger, we did not recover a high enough concentration for detection. Longer sampling times may also have been required, but due to

evaporation of the impingement solution this is often impossible, unless a non-evaporating fluid like mineral oil is used (Lin et al. 1999). In addition, it has been shown that bacteria can lose their ability to grow and form colonies on culturable plates due to physical stress of the sampling technique (Terzieva et al. 1996).

While PCR allows for the detection and identification of non-culturable airborne microorganisms, it does not allow one to distinguish between non-viable and viable microorganisms (Dungan and Leytem 2009b). Clones prepared from samples taken downwind from the cattle lots, lagoon, and compost pile were identified as being gram-negative and matched with the following genera: *Acinetobacter*, *Bradyrhizobium*, *Escherichia*, *Idiomarina*, *Methylobacterium*, *Ralstonia*, and *Novosphingobium* (Table 3). All of these genera were also identified in the upwind clones, except *Escherichia*, *Idiomarina*, and *Novosphingobium*. Downwind clones generally affiliated with gram-positive bacteria were identified as belonging to Firmicutes and Actinobacteria (Table 3). Immediately downwind from cattle feedlots, Wilson et al. (2002) found only non-pathogenic gram-positive bacteria such as *Bacillus*, *Chrysobacterium*, *Corynebacterium*, *Helococcus*, *Micrococcus*, and *Paenibacillus*. Since the majority of the clones in this study matched with sequences at the genus level or higher (not to species or strain level), knowledge of the pathogenicity of the airborne microorganisms could not be ascertained. In addition, due to limitations with DNA extraction and PCR amplification using universal primers, it should be noted that our survey is not comprehensive or even representative of all airborne bacterial genera and species. While many pathogenic strains of bacteria (e.g., *E. coli* and *Salmonella* spp.) are generally not considered a direct inhalation hazard, they are a potential infectious hazard when deposited in the throat and upper airway and swallowed (Pillai and Ricke 2002).

With respect to airborne coliphage as an indicator organism of viral contamination, the lack of positive recovery at the downwind sites could mean that the airborne concentrations were either too low and/or survivability of the coliphage in the aerosolized state is poor. In a study conducted by Carducci et al. (1999) at a sewage treatment plant, the airborne coliphage concentration decreased by 99% at 50 m, whereas reovirus and enterovirus only decreased by

15% at the same distance, suggesting that coliphage are not a good indicator of viral contamination. Results from the Carducci et al. (1999) study also found a significant correlation with total bacteria and fecal streptococci and aerosolized virus, possibly being more useful as an indicator of viral contamination than coliphage counts. The fact that we did not recover aerosolized coliphage in the downwind sites does not mean that virus, pathogenic or non-pathogenic, are not present. If total airborne bacterial counts could be used as an indicator of viral contamination, then one can speculate that the downwind samples contained virus as a result of elevated concentrations of airborne heterotrophs.

As with the heterotrophic bacteria, airborne fungal concentrations increased significantly at the downwind edge of the lot; however, the concentration did not decrease at 200 m downwind (Fig. 3). Compared to the upwind site with an average concentration of 515 CFU m<sup>-3</sup>, at DW1 and DW2, the concentrations were on average 1.8- and 2-fold higher, respectively. Regardless of lot conditions (Table 1), the airborne concentrations were highly variable during the course of the study (Table 4), despite the fact that lots were frozen and/or covered with snow in December, January, and February. Snow has been shown to reduce airborne concentrations by removing spores from the air and by covering ground sources (Li and Kendrick 1995). Fungal spores can be aerosolized through natural ejection mechanisms, but wind is often necessary for aerosolization and carrying the spores downwind (Jones and Harrison 2004). Rain is also important, as spore concentrations are higher after a rain event, as it helps release spores that resist dislodgment by wind. Additionally, air temperature, relative humidity, solar radiation, and wind speed have been shown to be positively and negatively correlated with airborne concentrations of various species of fungi (Jones and Harrison 2004). In our study, the total airborne fungal concentrations were also positively and negatively correlated with the ambient weather data, but not consistently at all sites. For example, air temperature at the upwind site was positively correlated with the airborne fungal concentrations, while at DW1 and DW2, it was negatively correlated. This is likely related to the source of the fungal spores between the two sites, differences in vegetative cover (i.e., surrounding land is irrigated crop land and lots are bare soil), and/or the

fact that correlations were based on average ambient weather data for the sampling period and did not consider diurnal effects.

In a study of airborne fungi in an indoor dairy cattle shed, fungal species from the following genera were recovered: *Absidia*, *Alternaria*, *Aspergillus*, *Cladosporium*, *Fusarium*, *Penicillium*, *Mucor*, *Rhizopus*, and *Syncephalastrum* (Adhikari et al. 2004). The concentration of culturable fungi at an outdoor station averaged about 500 CFU m<sup>-3</sup> and in the cattle sheds it was reported to range from 165 to 2,225 m<sup>-3</sup>, which is markedly similar to results obtained in our study at the upwind and downwind sites. While we did not determine the percent contribution of each fungal genera to the total culturable numbers, Adhikari et al. (2004) reported that more than 50% of the total fungal spores were contributed by *Aspergillus*, *Cladosporium*, *Nigrospora*, and *Penicillium*. In open-air swine houses, *Cladosporium* represented more than 90% of identified fungi (Chang et al. 2001). Other airborne fungal genera identified by Chang et al. (2001) were as follows: *Alternaria*, *Aspergillus* (*A. niger* and *flavus*), *Aureobasidium*, *Botrytis*, *Candida*, *Curvularia*, *Drechslera*, *Fusarium*, *Geotrichum*, *Monilia*, *Oidium*, *Paecilomyces*, *Penicillium*, *Sclerotium*, *Stemphylium*, *Trichoderma*, and *Ulocladium*. While some species within the genera identified in our study (Table 5) are known plant pathogens and can cause disease in immuno-compromised individuals, none of the fungal isolates are considered to be highly pathogenic to otherwise healthy humans (St-Germain and Summerbell 1996). However, it should be made clear that our fungal survey was not comprehensive or even representative of all airborne fungal genera and species.

## 5 Conclusions

The airborne heterotrophic bacterial concentrations were shown to fluctuate throughout the year-long study, with no correlation to ambient weather conditions. While total concentrations were higher immediately at the downwind edge of the lot, concentrations declined 200 m further downwind. A decrease in the bioaerosol concentration with distance from the source is a trend commonly seen in other similar studies. Even though the heterotrophic bacteria were not necessarily pathogenic microorganisms, this trend of decreasing concentration with distance from the source

is indicative of the transport behavior of pathogenic and non-pathogenic bacteria alike. Based upon the survey of a small number of bacterial clones from downwind sites, none of the sequence matches appeared to be affiliated with genera that contain species known to be pathogenic to humans. Airborne fungal concentrations at the open-lot dairy were also shown to increase at the edge of the lot, but stayed elevated at 200 m downwind. This is likely due to the fact that fungal spores are not associated with particulate matter, therefore can be carried farther downwind as a result of differing physical properties from particulate-bound bioaerosols. While a comprehensive survey of airborne fungi was not conducted, none of the fungal species identified in this study are known human pathogens, except in cases where patients may be immuno-compromised. Based upon our results, it is evident that decreasing ones proximity to the downwind perimeter of a CAFO, such as an open-lot dairy, will decrease exposure to airborne microorganisms. Reduced exposure could certainly translate into decreased potential for infection if pathogenic microbes are present at elevated concentrations. However, since epidemiological studies were not simultaneously conducted, the actual risk of infection at each downwind site cannot be determined from our results.

**Acknowledgments** We would like to thank numerous individuals at the dairy for accommodating us during this long study. We also thank the following individuals: Susie Hansen, Myles Miller, Patsy Heinemann, and Ann Pool.

## References

- Adhikari, A., Sen, M. M., Gupta-Bhattacharya, S., & Chanda, S. (2004). Volumetric assessment of airborne fungi in two sections of a rural indoor dairy cattle shed. *Environment International*, *29*, 1071–1078.
- Blanchard, D. C., & Syzdek, L. (1970). Mechanism for the water-to-air transfer and concentration of bacteria. *Science*, *170*, 626–628.
- Brandi, G., Sisti, M., & Amagliani, G. (2000). Evaluation of the environmental impact of microbial aerosols generated by wastewater treatment plants utilizing different aeration systems. *Journal of Applied Microbiology*, *88*, 845–852.
- Brooks, J. P., Gerba, C. P., & Pepper, I. L. (2004). Biological aerosol emission, fate, and transport from municipal and animal wastes. *Journal of Residuals Science & Technology*, *1*, 15–28.
- Brooks, J. P., Tanner, B. D., Josephson, K. L., Gerba, C. P., Haas, C. N., & Pepper, I. L. (2005). A national study on the residential impact of biological aerosols from the land application of biosolids. *Journal of Applied Microbiology*, *99*, 310–322.
- Brown, J. K. M., & Hovmoller, M. S. (2002). Aerial dispersal of pathogens on the global and continental scales and its impact on plant disease. *Science*, *297*, 537–541.
- Carducci, A., Gemelli, C., Cantiani, L., Casini, B., & Rovini, E. (1999). Assessment of microbial parameters as indicators of viral contamination of aerosol from urban sewage treatment plants. *Letters in Applied Microbiology*, *28*, 207–210.
- Centner, T. J. (2003). Regulating concentrated animal feeding operations to enhance the environment. *Environmental Science & Policy*, *6*, 433–440.
- Chang, C. W., Chung, H., Huang, C. F., & Su, H. J. J. (2001). Exposure of workers to airborne microorganisms in open-air swine houses. *Applied and Environmental Microbiology*, *67*, 155–161.
- Cox, C. S. (1995a). Physical aspects of bioaerosol particles. In C. S. Cox & C. M. Wathes (Eds.), *Bioaerosols Handbook* (pp. 15–25). New York: Lewis Publishers.
- Cox, C. S. (1995b). Stability of airborne microbes and allergens. In C. S. Cox & C. M. Wathes (Eds.), *Bioaerosols handbook* (pp. 77–99). New York: Lewis Publishers.
- Crook, B., & Sherwood-Higham, J. L. (1997). Sampling and assay of bioaerosols in the work environment. *Journal of Aerosol Science*, *28*, 417–426.
- Dungan, R. S., & Leytem, A. B. (2009a). Airborne endotoxin concentrations at a large open-lot dairy in southern Idaho. *Journal of Environmental Quality*, *38*, 1919–1923.
- Dungan, R. S., & Leytem, A. B. (2009b). Qualitative and quantitative methodologies for determination of airborne microorganisms at concentrated animal-feeding operations. *World Journal of Microbiology and Biotechnology*, *25*, 1505–1518.
- Eaton, A. D., Clesceri, L. S., Rice, E. W., & Greenberg, A. E. (2005). *Standard methods for the examination of water and wastewater* (21st ed.). Washington DC: American Public Health Association, American Water Works Association, Water Environment Federation.
- Ehrlich, R., Miller, S., & Walker, R. L. (1970). Relationship between atmospheric temperature and survival of airborne bacteria. *Applied Microbiology*, *19*, 245–249.
- Fracchia, L., Pietronave, S., Rinaldi, M., & Martinotti, M. G. (2006). The assessment of airborne bacterial contamination in three composting plants revealed site-related biological hazard and seasonal variations. *Journal of Applied Microbiology*, *100*, 973–984.
- Gardes, M., & Bruns, T. D. (1993). ITS primers with enhanced specificity for basidiomycetes—application to the identification of mycorrhizae and rusts. *Molecular Ecology*, *2*, 113–118.
- Green, C. F., Gibbs, S. G., Tarwater, P. M., Mota, L. C., & Scarpino, P. V. (2006). Bacterial plume emanating from the air surrounding swine confinement operations. *Journal of Occupational and Environmental Hygiene*, *3*, 9–15.
- Hall, T. A. (1999). BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, *41*, 95–98.
- Heederik, D., Sigsgaard, T., Thorne, P. S., Kline, J. N., Avery, R., Bønløkke, J. H., et al. (2007). Health effects of airborne

- exposures from concentrated animal feeding operations. *Environmental Health Perspectives*, 115, 298–302.
- Jones, A. M., & Harrison, R. M. (2004). The effects of meteorological factors on atmospheric bioaerosol concentrations—a review. *Science of the Total Environment*, 326, 151–180.
- Kiekhaefer, M. S., Donham, K. J., Whitten, P., & Thorne, P. S. (1995). Cross seasonal studies of airborne microbial populations and environment in swine buildings: Implications for worker and animal health. *Annals of Agricultural and Environmental Medicine*, 2, 37–44.
- Li, D.-W., & Kendrick, B. (1995). A year-round study of functional relationships of airborne fungi with meteorological factors. *International Journal of Biometeorology*, 39, 74–80.
- Lighthart, B. (1994). Physics of microbial bioaerosols. In B. Lighthart & A. J. Mohr (Eds.), *Atmospheric microbial aerosols: Theory and applications* (pp. 5–27). New York: Chapman & Hall.
- Lin, X., Reponen, T. A., Willeke, K., Grinshpun, S. A., Foarde, K. K., & Ensor, D. S. (1999). Long-term sampling of airborne bacteria and fungi into a non-evaporating liquid. *Atmospheric Environment*, 33, 4291–4298.
- Marchesi, J. R., Sato, T., Weightman, A. J., Martin, T. A., Fry, J. C., Hiom, S. J., et al. (1998). Design and evaluation of bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied and Environmental Microbiology*, 64, 795–799.
- Marthi, B., Fieland, V. P., Walter, M., & Seidler, R. J. (1990). Survival of bacteria during aerosolization. *Applied and Environmental Microbiology*, 56, 3463–3467.
- Millner, P. D. (2009). Bioaerosols associated with animal production operations. *Bioresource Technology*, 100, 5379–5385.
- Mohr, A. J. (2007). Fate and transport of microorganisms in air. In C. J. Hurst, et al. (Eds.), *Manual for environmental microbiology* (pp. 952–960). Washington DC: ASM Press.
- Muyzer, G., DeWaal, E. C., & Uitterlinden, A. G. (1993). Profiling complex microbial populations by denaturing gradient gel electrophoresis analysis by polymerase chain reaction-amplified genes coding for 16s rRNA. *Applied and Environmental Microbiology*, 59, 695–700.
- Nicholson, K. W. (1995). Physical aspects of bioaerosol sampling and deposition. In C. S. Cox & C. M. Wathes (Eds.), *Bioaerosols handbook* (pp. 27–53). New York: Lewis Publishers.
- Pillai, S. D., & Ricke, S. C. (2002). Bioaerosols from municipal and animal wastes: Background and contemporary issues. *Canadian Journal of Microbiology*, 48, 681–696.
- Poon, C. P. C. (1966). Studies on the instantaneous death of airborne *Escherichia coli*. *American Journal of Epidemiology*, 84, 1–9.
- Reysenback, A. L., Wickham, G. S., & Pace, N. R. (1994). Phylogenetic analysis of the hyperthermophilic pink filament community in Octopus Spring, Yellowstone National Park. *Applied and Environmental Microbiology*, 60, 2113–2119.
- Rule, A. M., Chapin, A. R., McCarthy, S. A., Gibson, K. E., Schwab, K. J., & Buckley, T. J. (2005). Assessment of an aerosol treatment to improve air quality in a swine concentrated animal feeding operation (CAFO). *Environmental Science and Technology*, 39, 9649–9655.
- SAS Institute. (2004). *SAS/STAT user's guide. Release 9.1*. Cary, NC: SAS Institute, Inc.
- Schulze, A., van Strien, R., Ehrenstein, V., Schierl, R., Küchenhoff, H., & Radon, K. (2006). Ambient endotoxin level in an area with intensive livestock production. *Annals of Agricultural and Environmental Medicine*, 13, 87–91.
- Songer, J. R. (1967). Influence of relative humidity on the survival of some airborne viruses. *Applied Microbiology*, 15, 35–42.
- Spaan, S., Wouters, I. M., Oosting, I., Doekes, G., & Heederik, D. (2006). Exposure to inhalable dust and endotoxins in agricultural industries. *Journal of Environmental Monitoring*, 8, 63–72.
- Stetzenbach, L. D. (2007). Introduction to aerobiology. In C. J. Hurst, et al. (Eds.), *Manual for environmental microbiology* (pp. 952–960). Washington, DC: ASM Press.
- St-Germain, G., & Summerbell, R. (1996). *Identifying filamentous fungi: A clinical laboratory handbook*. Belmont, CA: Star Publishing Co.
- Sweeten, J. M., Parnell, C. B., Etheredge, R. S., & Osborne, D. (1988). Dust emissions in cattle feedlots. *Veterinary Clinics of North America. Food Animal Practice*, 4, 557–578.
- Taha, M. P. M., Pollard, S. J. T., Sarkar, U., & Longhurst, P. (2005). Estimating fugitive bioaerosol releases from static compost windrows: Feasibility of a portable wind tunnel approach. *Waste Management*, 25, 445–450.
- Terzieva, S., Donnelly, J., Ulevicius, V., Grinshpun, S. A., Willeke, K., Stelma, G. N., et al. (1996). Comparison of methods for detection and enumeration of airborne microorganisms collected by liquid impingement. *Applied and Environmental Microbiology*, 62, 2264–2272.
- US Environmental Protection Agency. (2002). *Method 1604: Total Coliforms and Escherichia coli in water by membrane filtration using a simultaneous detection technique (MI medium)*. EPA 821-R-02-024. Washington, DC: Office of Water.
- USDA, National Agricultural Statistics Service [online]. (2008). Available at [www.nass.usda.gov](http://www.nass.usda.gov) (verified on 2 June 2008).
- White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. A. Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR protocols: A guide to methods and applications* (pp. 315–322). New York: Academic Press.
- Wickman, H. H. (1994). Deposition, adhesion, and release. In B. Lighthart & A. J. Mohr (Eds.), *Atmospheric microbial aerosols*. New York: Chapman & Hall.
- Wilson, S. C., Morrow-Tesch, J., Straus, D. C., Cooley, J. D., Wong, W. C., Mitlohner, F. M., et al. (2002). Airborne microbial flora in a cattle feedlot. *Applied and Environmental Microbiology*, 68, 3238–3242.
- Zucker, B.-A., & Müller, W. (1998). Concentrations of airborne endotoxin in cow and calf stables. *Journal of Aerosol Science*, 29, 217–221.
- Zucker, B.-A., Trojan, S., & Müller, W. (2000). Airborne gram-negative bacterial flora in animal houses. *Journal of Veterinary Medicine, B*, 47, 37–46.